

POSTER PRESENTATION

Open Access

Ion concentration dynamics leads to the very slow spontaneous neuronal oscillations

Giri P Krishnan, Oscar Gonzales, Maxim Bazhenov*

From The Twenty Third Annual Computational Neuroscience Meeting: CNS*2014
Québec City, Canada. 26-31 July 2014

Resting or baseline brain activity may fluctuate at the very low frequencies (0.01-0.1 Hz) [1]. These oscillations, revealed by fMRI data, are found to be correlated across brain regions. There has been rising interest to this slow activity, since it may reflect functional connectivity between brain regions and strongly correlate with neuronal firing.

In this study we propose that spontaneous slow brain oscillations may rise from ionic concentration dynamics. We used biophysically realistic computational model of the cortical network based on the Hodgkin-Huxley kinetics and implementing ion concentration dynamics. Computational network model consisted of pyramidal and inhibitory neurons with connectivity based on AMPA, GABA_A and NMDA synapses. It implemented dynamic variables for intra- and extracellular K⁺/Na⁺ concentrations, intracellular Cl⁻ concentration and simulated Na⁺/K⁺ exchange and KCC2 pumps. Random Poisson input was given to each neuron to mimic in vivo conditions.

Our study revealed that extracellular K⁺ dynamics determine the slow network oscillations in the frequency range 0.01-0.05 Hz. The network oscillations were more prominent for larger network compared to the small networks. The oscillation amplitude depended on the random input to the network. While K⁺ dynamics and diffusion contributed to the local synchrony, long-range synaptic connectivity provided correlated activity between distinct populations of neurons with no ion diffusion between sites. This finding may explain observation that structural network connections determine the functional connectivity in human recordings [2]. We conclude that the ion concentration dynamics may contribute to the slow spontaneous fluctuations in brain activity found with fMRI

recordings. Future experiments are required to verify this prediction.

Published: 21 July 2014

References

1. Fox MD, Raichle ME: Spontaneous fluctuations in brain activity observed with functional magnetic resonance imaging. *Nature Reviews Neuroscience* 2007, **8**(9):700-711.
2. Honey CJ, Kötter R, Breakspear M, Sporns O: Network structure of cerebral cortex shapes functional connectivity on multiple time scales. *Proceedings of the National Academy of Sciences* 2007, **104**(24):10240-10245.

doi:10.1186/1471-2202-15-S1-P217

Cite this article as: Krishnan et al.: Ion concentration dynamics leads to the very slow spontaneous neuronal oscillations. *BMC Neuroscience* 2014 **15**(Suppl 1):P217.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit



* Correspondence: maxim.bazhenov@ucr.edu
Department of Cell Biology & Neuroscience, University of California,
Riverside, CA, USA